

Compressibility of electron gas in presence of Anderson's impurity.

Daniel L. Miller

Dept. of Physics of Complex Systems,
The Weizmann Institute of science, Rehovot, 76100 Israel
e-mail fndaniil@wicc.weizmann.ac.il

(February 1, 2008)

We employ Wiegmann's solution of the Anderson impurity model in order to compute the compressibility of electron gas. We have found that there is a pair of neighbor levels separated by anomalously large energy $\propto L^{-1/3}$, where L is the size of the system. The Fermi energy crosses this pseudogap and the compressibility decreases near the so-called "symmetric" limit.

The Anderson Hamiltonian¹ describes the localized state interacting with electron gas

$$\hat{\mathcal{H}} = \sum_{k\sigma} \left\{ \epsilon_k \hat{a}_{k\sigma}^\dagger \hat{a}_{k\sigma} + V_k \hat{a}_{k\sigma}^\dagger \hat{a}_{d\sigma} + V_k^* \hat{a}_{d\sigma}^\dagger \hat{a}_{k\sigma} \right\} + \epsilon_d (\hat{n}_\uparrow + \hat{n}_\downarrow) + U \hat{n}_\uparrow \hat{n}_\downarrow \quad \hat{n}_\sigma = \hat{a}_{d\sigma}^\dagger \hat{a}_{d\sigma} . \quad (1)$$

Here $\hat{a}_{k\sigma}^\dagger$ creates the electron in the band with wave number k , spin σ , and energy ϵ_k . The operator $\hat{a}_{d\sigma}^\dagger$ creates the localized electron with energy ϵ_d . The interaction between localized electrons gives the correlation term $U \hat{n}_\uparrow \hat{n}_\downarrow$, and $V_k \hat{a}_{k\sigma}^\dagger \hat{a}_{d\sigma}$ describes their decay to the band.

Wiegmann have solved this problem in one dimension². At the zero magnetic field the ground state is described by the set of rapidities $\{\lambda_\alpha\}_{\alpha=0}^M$. They satisfy

$$\pi J_\alpha + \sum_{\beta=0}^M \arctan(\lambda_\alpha - \lambda_\beta) = -x(\lambda_\alpha) L \quad (2)$$

$$x(\lambda) = \epsilon_d + U/2 - \sqrt{UT(\lambda + \sqrt{\lambda^2 + 1/4})}, \quad (3)$$

where L is the size of the sample, $\Gamma = |V|^2$, $\hbar v_F = 1$, and therefore $\epsilon_k = k$. Terms of the order of $1/L$ are omitted, because we are going to discuss the properties of the electron gas only. The integer numbers $\{J_\alpha\}_{\alpha=0}^M$ completely describe the state of the system with $N = 2(M+1)$ particles. The energy of the system is just

$$E = 2 \sum_{\alpha=0}^M x(\lambda_\alpha). \quad (4)$$

The ground state corresponds to subsequent values of J_α ; we can put $J_\alpha = \frac{LD}{\pi} - \frac{3M}{2} \dots \frac{LD}{\pi} + \frac{M}{2}$, where D is the energy of the bottom of the band. In the non-interacting case, $U = 0$, one obtains the chemical potential $\mu \equiv \frac{dE}{dN} = \frac{2\pi M}{L} - D$. The compressibility of the gas is $\frac{dn}{d\mu} = \frac{1}{\pi}$, where $n = N/L$ is the density of particles.

These results remain valid for finite U if one makes use of the following continuous approximation. Under the condition

$$|\lambda_{\alpha+1} - \lambda_\alpha| \ll 1 \quad \forall \alpha \quad (5)$$

one can transform Eq. (2) to the integral equation

$$\sigma_\lambda + \tilde{\sigma}_\lambda + \frac{1}{\pi} \int_{\lambda_0}^{\lambda_M} \frac{\sigma_{\lambda'} d\lambda'}{1 + (\lambda - \lambda')^2} = -\frac{1}{\pi} \frac{dx}{d\lambda} \quad (6)$$

where the mean density of rapidities $\sigma(\lambda) = \frac{1/L}{\lambda_{\alpha+1} - \lambda_\alpha}$, and $\sigma(\lambda) \propto \theta(\lambda - \lambda_0)$. (Step function.) We have also introduced the auxiliary function $\tilde{\sigma}(\lambda) \propto \theta(\lambda_0 - \lambda)$ in order to make Eq. (6) valid for $\lambda \in]-\infty, \lambda_M]$.

It is convenient to integrate Eq. (6) from $-\infty$ to λ_M and combine it with Eq. (2) for $\alpha = 0$

$$\pi \int_{-\infty}^{\lambda_0} \tilde{\sigma}_\lambda d\lambda = \epsilon_d + U/2 + D - 2\pi \frac{M}{L}. \quad (7)$$

This is an equation for λ_0 . In the symmetric limit $|\epsilon_d + U/2 + D - 2\pi \frac{M}{L}| \ll \sqrt{UT}$ one has³

$$\lambda_0 \approx \frac{1}{\pi} \log \frac{E_d + U/2 + D - 2\pi M/L}{2\sqrt{UT/\pi e}} \quad (8)$$

goes to $-\infty$ when $E_d + U/2 + D - 2\pi M/L$ goes to zero.

The problem is that the condition Eq. (5) will be broken eventually. It is clear that for $\lambda_0 \rightarrow -\infty$ the density of the rapidities goes to zero $\sigma(\lambda_0) < |\frac{dx}{d\lambda}| \sim |\lambda_0|^{-3/2}$. Therefore

$$|\lambda_1 - \lambda_0| \gtrsim |\lambda_0|^{3/2} \rightarrow \infty, \quad (9)$$

and the solution of Eq. (2) cannot be described by the density of rapidities $\sigma(\lambda)$.

The main idea of the present paper is to introduce one rapidity λ_0 separated from all the rest $\lambda_1 \dots \lambda_M$. Then

$$x(\lambda_0) = 2\pi \frac{M}{L} - D \quad \lambda_0 \approx -\frac{UT/8}{(2\pi M/L - D - \epsilon_d - U/2)^2} \quad (10)$$

is obtained directly from Eq. (2) with $\alpha = 0$ under the condition $\lambda_\alpha - \lambda_0 \gg 1$ for $\alpha = 1 \dots M$. The rest of equations gives

$$\sigma_\lambda + \tilde{\sigma}_\lambda + \frac{1}{\pi} \int_{\lambda_1}^{\lambda_M} \frac{\sigma_{\lambda'} d\lambda'}{1 + (\lambda - \lambda')^2} = -\frac{1}{\pi} \frac{dx}{d\lambda} \quad (11)$$




FIG. 1. The main parameters of the problem. Energy levels appear densely between $-D$ and $\mu \sim \epsilon_d + U/2$ (“symmetric” limit). Let us just mention “asymmetric” limit, when Fermi level is near ϵ_d ; this is the regime of the formation of the local moment.

The solution is well known and instead of Eq. (8) we will have

$$\lambda_1 \approx \frac{1}{2\pi} \log \frac{1}{|\lambda_0|} \quad (12)$$

It means appearance of a gap in the excitation spectrum and a jump of the chemical potential

$$\delta\mu = 2[x(\lambda_0) - x(\lambda_1)] \approx \sqrt{\frac{\pi U \Gamma}{\log |\lambda_0|}}. \quad (13)$$

The gap is opened when the condition Eq. (5) is broken for $\lambda_1 - \lambda_0$. This occurs when $(L/\pi)|dx/d\lambda| \approx 1$ or

$$2\lambda_1 \approx \left(\frac{U \Gamma L^2}{4\pi^2} \right)^{1/3} \quad (14)$$

So our main result is derived from Eqs. (12)-(14)

$$\delta\mu \approx \left(\frac{\pi U \Gamma}{L} \right)^{1/3} \quad (15)$$

and the gap is anomalously large, because it is proportional to $L^{-1/3}$. The compressibility is suppressed $\frac{\delta n}{\delta \mu} \sim (U \Gamma L^2)^{-1/3} \ll 1$.

It is important to mention that the model above is valid for even number of electrons in the system, and the computed above gap is for two additional particles.

The energy of the intermediate state with odd number of electrons can be computed also by the method presented here. It is necessarily to introduce equation for the charge rapidity and to couple it to Eq. (2). Then our huge gap will be split in two parts. However, at least one of them must be large.

In conclusion we have found the anomalously large spacing between energy levels of electron gas interacting with impurity. This is a finite size effect, which requires special treatment. The pseudogap appears at the Fermi energy when the model approaches complete particle-hole symmetry, $\mu \approx \epsilon_d + U/2$.

ACKNOWLEDGMENTS

This work was supported by Israel Science Foundation and the Minerva Center for Nonlinear Physics of Complex systems.

¹ P. W. Anderson, Phys. Rev. **124**, 41 (1961).

² P. B. Wiegmann, Phys. Lett. A **80**, 163 (1980).

³ P. B. Wiegmann, J. Phys. C **16**, 2281 (1983).